

Homework #3

Due: Friday February 20

(50 pts)

MTHBD/CMPBD 424

1. Suppose that a differential equation is solved numerically on an interval $[a, b]$ and at each step you get a local truncation error $= kh^p$ where h is the step size $= \frac{b-a}{N}$ and $k > 0$. Assuming that the only error is this local truncation error at each step, show that the total local truncation error $= (b - a)kh^{p-1}$.

We showed this for the case of $p = 2$ in class. We also did a much more complicated proof to show that this method had a global discretization error $= \mathcal{O}(h)$. I am asking for the simpler case only with p arbitrary.

2. Consider the initial value problem

$$x' = x + t \quad x(0) = 0$$

We numerically solved this problem in class using Euler's method and the Taylor series method of order 2. I want you to solve this problem using Euler's method and the Taylor series method of order 3. The exact solution is $x = e^t - (t + 1)$. I want you to find the numerical solution for $0 \leq t \leq 2$.

Hand in

- (a) A page describing the steps in the Taylor series method of order 3 for this particular problem.
- (b) A graph of the exact solution, Euler's method and the Taylor series method of order 3 all on the same graph for $0 \leq t \leq 2$. For these let $h = .1$.
- (c) Create a vector of step sizes $h = [2^{-3}, 2^{-4}, \dots, 2^{-10}]$. Let E_h represent the absolute value of the error at $t = 2$ between the exact solution and Euler's method for a given value of h . Hand in a graph of $\log_{10}(E_h)$ versus $\log_{10}(h)$ over the complete range of h values. Be sure to put $\log_{10}(h)$ on the x-axis and $\log_{10}(E_h)$ on the y-axis of your graph. On the same page comment on the slope of this curve and what this implies about the error of Euler's method.
- (d) Do the same as part (C) for the Taylor Series Method of Order 3.

1. Suppose that a differential equation is solved numerically on an interval $[a, b]$ and at each step you get a local truncation error $= kh^p$ where h is the step size $= \frac{b-a}{N}$ and $k > 0$. Assuming that the only error is this local truncation error at each step, show that the total local truncation error $= (b - a)kh^{p-1}$.

Hand In:

- **Page 1:** This page with the answer to problem 1.
- **Page 2:** The answer to problem 2(a).
- **Page 3:** The graph from problem 2(b).
- **Page 4:** The graph and comments from problem 2(c).
- **Page 5:** The graph and comments from problem 2(d).
- **Page \geq 6:** The code for your Taylor series method of order 3.